

**A new approach to the series expansion of iterated Stratonovich stochastic integrals of arbitrary multiplicity with respect to components of the multidimensional Wiener process**

**Dmitriy F. Kuznetsov**

Peter the Great Saint-Petersburg Polytechnic University, Russia

E-mail: sde\_kuznetsov@inbox.ru

The work is devoted to a new approach to the series expansion of iterated Stratonovich stochastic integrals with respect to components of the multidimensional Wiener process. This approach is based on multiple Fourier–Legendre series as well as multiple trigonometric Fourier series. The theorem on the mean-square convergent expansion for the iterated Stratonovich stochastic integrals of arbitrary multiplicity is formulated and proved under the condition of convergence of trace series. This condition has been verified for integrals of multiplicities 2 to 5 and complete orthonormal systems of Legendre polynomials and trigonometric functions in Hilbert space. The Hu–Meyer formula [1] and multiple Wiener–Itô stochastic integral [2] were used in the proof of the mentioned theorem. The rate of mean-square convergence of the obtained expansions is found. The mentioned results can be applied to the numerical integration of Itô stochastic differential equations with non-commutative noise in the framework of the approach based on the Taylor–Stratonovich expansion.

Let  $(\Omega, \mathbb{F}, \mathbb{P})$  be a complete probability space, let  $\{\mathbb{F}_\tau, \tau \in [0, T]\}$  be a nondecreasing right-continuous family of  $\sigma$ -algebras of  $\mathbb{F}$ , and let  $\mathbf{W}_\tau$  be a standard  $m$ -dimensional Wiener stochastic process, which is  $\mathbb{F}_\tau$ -measurable for any  $\tau \in [0, T]$ . We assume that the components  $\mathbf{W}_\tau^{(i)}$  ( $i = 1, \dots, m$ ) of this process are independent.

Consider the unordered set  $\{1, 2, \dots, k\}$  and separate it into two parts: the first part consists of  $r$  unordered pairs (sequence order of these pairs is also unimportant) and the second one consists of the remaining  $k - 2r$  numbers. So, we have

$$(1) \quad \left( \underbrace{\{\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}\}}_{\text{part 1}}, \underbrace{\{q_1, \dots, q_{k-2r}\}}_{\text{part 2}} \right),$$

where  $\{g_1, g_2, \dots, g_{2r-1}, g_{2r}, q_1, \dots, q_{k-2r}\} = \{1, 2, \dots, k\}$ , braces mean an unordered set, and parentheses mean an ordered set.

Consider the Fourier coefficient

$$(2) \quad C_{j_k \dots j_1} = \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \dots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \dots dt_k$$

corresponding to the function  $K(t_1, \dots, t_k) = \psi_1(t_1) \dots \psi_k(t_k) \mathbf{1}_{\{t_1 < \dots < t_k\}}$ ,  $t_1, \dots, t_k \in [t, T]$ ,  $k \geq 2$  and  $K(t_1) = \psi_1(t_1)$ ,  $t_1 \in [t, T]$ . Here  $\psi_1(\tau), \dots, \psi_k(\tau) : [t, T] \rightarrow \mathbb{R}$  are nonrandom functions,  $\{\phi_j(x)\}_{j=0}^\infty$  is a complete orthonormal system of functions in the space  $L_2([t, T])$ ,  $j_1, \dots, j_k = 0, 1, \dots$ ,  $\mathbf{1}_A$  denotes the indicator of the set  $A$ . At that we suppose  $\phi_0(x) = 1/\sqrt{T-t}$ . Here and further, we assume that  $0 \leq t < T < \infty$ .

Denote

$$(3) \quad C_{j_k \dots j_{l+1} j_l j_{l-2} \dots j_1} \Big|_{(j_l j_l) \sim (\cdot)} \stackrel{\text{def}}{=} \int_t^T \psi_k(t_k) \phi_{j_k}(t_k) \dots \int_t^{t_{l+2}} \psi_{l+1}(t_{l+1}) \phi_{j_{l+1}}(t_{l+1}) \int_t^{t_{l+1}} \psi_l(t_l) \psi_{l-1}(t_l) \times \\ \times \int_t^{t_l} \psi_{l-2}(t_{l-2}) \phi_{j_{l-2}}(t_{l-2}) \dots \int_t^{t_2} \psi_1(t_1) \phi_{j_1}(t_1) dt_1 \dots dt_{l-2} dt_l t_{l+1} \dots dt_k,$$

i.e.  $C_{j_k \dots j_{l+1} j_l j_{l-2} \dots j_1} \Big|_{(j_l j_l) \sim (\cdot)}$  is again the Fourier coefficient of type  $C_{j_k \dots j_1}$  but with a new shorter multi-index  $j_k \dots j_{l+1} 0 j_{l-2} \dots j_1$  and new weight functions  $\psi_1(\tau), \dots, \psi_{l-2}(\tau), \sqrt{T-t} \psi_{l-1}(\tau) \psi_l(\tau)$ ,

$\psi_{l+1}(\tau), \dots, \psi_k(\tau)$  (also we suppose that  $\{l, l-1\}$  is one of the pairs  $\{g_1, g_2\}, \dots, \{g_{2r-1}, g_{2r}\}$  from the relation (1)).

Denote

$$\begin{aligned} \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} &\stackrel{\text{def}}{=} \sum_{j_{g_{2r-1}}=p+1}^{\infty} \sum_{j_{g_{2r-3}}=p+1}^{\infty} \dots \sum_{j_{g_3}=p+1}^{\infty} \sum_{j_{g_1}=p+1}^{\infty} C_{j_k \dots j_1} \Big|_{j_{g_1}=j_{g_2}, \dots, j_{g_{2r-1}}=j_{g_{2r}}}, \\ S_l \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} \right\} &\stackrel{\text{def}}{=} \frac{1}{2} \mathbf{1}_{\{g_{2l}=g_{2l-1}+1\}} \sum_{j_{g_{2r-1}}=p+1}^{\infty} \sum_{j_{g_{2r-3}}=p+1}^{\infty} \dots \\ \dots \sum_{j_{g_{2l+1}}=p+1}^{\infty} \sum_{j_{g_{2l-3}}=p+1}^{\infty} \dots \sum_{j_{g_3}=p+1}^{\infty} \sum_{j_{g_1}=p+1}^{\infty} C_{j_k \dots j_1} &\Big|_{(j_{g_{2l}} j_{g_{2l-1}}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, \dots, j_{g_{2r-1}}=j_{g_{2r}}}. \end{aligned}$$

Note that the operation  $S_l$  ( $l = 1, 2, \dots, r$ ) acts on the value  $\bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}}$  as follows:  $S_l$  multiplies  $\bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}}$  by  $\mathbf{1}_{\{g_{2l}=g_{2l-1}+1\}}/2$ , removes the summation with respect to  $j_{g_{2l-1}}$ , and replaces  $C_{j_k \dots j_1} \Big|_{j_{g_1}=j_{g_2}, \dots, j_{g_{2r-1}}=j_{g_{2r}}}$  with  $C_{j_k \dots j_1} \Big|_{(j_{g_{2l}} j_{g_{2l-1}}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, \dots, j_{g_{2r-1}}=j_{g_{2r}}}$ . At that we write  $C_{j_k \dots j_1} \Big|_{(j_{g_1} j_{g_2}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}} = C_{j_k \dots j_1} \Big|_{(j_{g_1} j_{g_1}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}}$ .

The action of superposition  $S_l S_m$  on  $\bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}}$  is obvious. For example, for  $r = 3$

$$\begin{aligned} S_3 S_1 \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_5, g_6} \right\} &= \\ = \frac{1}{2^2} \mathbf{1}_{\{g_6=g_5+1\}} \mathbf{1}_{\{g_2=g_1+1\}} \sum_{j_{g_3}=p+1}^{\infty} C_{j_k \dots j_1} &\Big|_{(j_{g_2} j_{g_1}) \curvearrowright (\cdot), (j_{g_6} j_{g_5}) \curvearrowright (\cdot), j_{g_1}=j_{g_2}, j_{g_3}=j_{g_4}, j_{g_5}=j_{g_6}}. \end{aligned}$$

**Theorem 1** [3] (Sect. 2.10). *Assume that the continuously differentiable functions  $\psi_l(\tau)$  ( $l = 1, \dots, k$ ) and the complete orthonormal system  $\{\phi_j(x)\}_{j=0}^{\infty}$  of continuous functions ( $\phi_0(x) = 1/\sqrt{T-t}$ ) in the space  $L_2([t, T])$  are such that the following conditions are satisfied:*

1. *The equality*

$$(4) \quad \frac{1}{2} \int_t^s \Phi_1(\tau) \Phi_2(\tau) d\tau = \sum_{j=0}^{\infty} \int_t^s \Phi_2(\tau) \phi_j(\tau) \int_t^{\tau} \Phi_1(\theta) \phi_j(\theta) d\theta d\tau$$

*holds for all  $s \in (t, T]$ , where the nonrandom functions  $\Phi_1(\tau), \Phi_2(\tau)$  are continuously differentiable on  $[t, T]$  and the series on the right-hand side of (4) converges absolutely.*

2. *The estimates*

$$\left| \int_t^s \phi_j(\tau) \Phi_1(\tau) d\tau \right| + \left| \int_s^T \phi_j(\tau) \Phi_2(\tau) d\tau \right| \leq \frac{\Psi_1(s)}{j^{1/2+\alpha}}, \quad \left| \sum_{j=p+1}^{\infty} \int_t^s \Phi_2(\tau) \phi_j(\tau) \int_t^{\tau} \Phi_1(\theta) \phi_j(\theta) d\theta d\tau \right| \leq \frac{\Psi_2(s)}{p^\beta}$$

*hold for all  $s \in (t, T)$  and for some  $\alpha, \beta > 0$ , where  $\Phi_1(\tau), \Phi_2(\tau)$  are continuously differentiable nonrandom functions on  $[t, T]$ ,  $j, p \in \mathbb{N}$ , and*

$$\int_t^T \Psi_1^2(\tau) d\tau < \infty, \quad \int_t^T |\Psi_2(\tau)| d\tau < \infty.$$

3. *The condition*

$$\lim_{p \rightarrow \infty} \sum_{\substack{j_1, \dots, j_q, \dots, j_k=0 \\ q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}}}^p \left( S_{l_1} S_{l_2} \dots S_{l_d} \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} \right\} \right)^2 = 0$$

holds for all possible  $g_1, g_2, \dots, g_{2r-1}, g_{2r}$  (see (1)) and  $l_1, l_2, \dots, l_d$  such that  $l_1, l_2, \dots, l_d \in \{1, 2, \dots, r\}$ ,  $l_1 > l_2 > \dots > l_d$ ,  $d = 0, 1, 2, \dots, r-1$ , where  $r = 1, 2, \dots, [k/2]$  and

$$S_{l_1} S_{l_2} \dots S_{l_d} \left\{ \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} \right\} \stackrel{\text{def}}{=} \bar{C}_{j_k \dots j_q \dots j_1}^{(p)} \Big|_{q \neq g_1, g_2, \dots, g_{2r-1}, g_{2r}} \quad \text{for } d = 0.$$

Then, for the iterated Stratonovich stochastic integral of arbitrary multiplicity  $k$

$$(5) \quad J^*[\psi^{(k)}]_{T,t}^{(i_1 \dots i_k)} = \int_t^T \psi_k(t_k) \dots \int_t^{t_2} \psi_1(t_1) \circ d\mathbf{W}_{t_1}^{(i_1)} \dots \circ d\mathbf{W}_{t_k}^{(i_k)}$$

the following expansion

$$(6) \quad J^*[\psi^{(k)}]_{T,t}^{(i_1 \dots i_k)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_k=0}^p C_{j_k \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_k}^{(i_k)}$$

that converges in the mean-square sense is valid, where  $C_{j_k \dots j_1}$  is the Fourier coefficient defined by (2), l.i.m. is a limit in the mean-square sense,  $i_1, \dots, i_k = 0, 1, \dots, m$ ,

$$\zeta_j^{(i)} = \int_t^T \phi_j(\tau) d\mathbf{W}_\tau^{(i)}$$

are independent standard Gaussian random variables for various  $i$  or  $j$  (in the case when  $i \neq 0$ ),  $d\mathbf{W}_\tau^{(i)}$  and  $\circ d\mathbf{W}_\tau^{(i)}$  ( $i = 0, 1, \dots, m$ ) are Itô and Stratonovich differentials, respectively;  $\mathbf{W}_\tau^{(0)} \stackrel{\text{def}}{=} \tau$ .

The following theorem is proved [3] on the base of Theorem 1.

**Theorem 2** [3] (Sect. 2.11–2.15). Suppose that  $\{\phi_j(x)\}_{j=0}^\infty$  is a complete orthonormal system of Legendre polynomials or trigonometric functions in the space  $L_2([t, T])$ . Furthermore, let  $\psi_1(\tau), \dots, \psi_5(\tau)$  are continuously differentiable nonrandom functions on  $[t, T]$ . Then, for the iterated Stratonovich stochastic integral  $J^*[\psi^{(k)}]_{T,t}^{(i_1 \dots i_k)}$  ( $k = 3, 4, 5$ ) defined by (5) the following relations

$$(7) \quad J^*[\psi^{(k)}]_{T,t}^{(i_1 \dots i_k)} = \text{l.i.m.}_{p \rightarrow \infty} \sum_{j_1, \dots, j_k=0}^p C_{j_k \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_k}^{(i_k)},$$

$$(8) \quad \mathbf{M} \left( J^*[\psi^{(k)}]_{T,t}^{(i_1 \dots i_k)} - \sum_{j_1, \dots, j_k=0}^p C_{j_k \dots j_1} \zeta_{j_1}^{(i_1)} \dots \zeta_{j_k}^{(i_k)} \right)^2 \leq \frac{C}{p^{1-\varepsilon}}$$

are fulfilled, where  $i_1, \dots, i_k = 0, 1, \dots, m$  in (7) and  $i_1, \dots, i_k = 1, \dots, m$  in (8), constant  $C$  is independent of  $p$ ,  $\varepsilon$  is an arbitrary small positive real number for the case  $k = 4, 5$  (polynomial case) and  $\varepsilon = 0$  for the case  $k = 3$  (polynomial and trigonometric cases) or for the case  $k = 4, 5$  (trigonometric case),  $C_{j_k \dots j_1}$  is the Fourier coefficient defined by (2), another notations are the same as in Theorem 1.

Note that (7), (8) are also valid for the case  $k = 1$  with  $\varepsilon = 0$  (see Theorems 1.1, 1.16 and Remark 1.7 [3]). The case  $k = 2$  of Theorem 2 as well as some narrow special cases of Theorem 2 for  $k = 3, 4$  were considered earlier in [3] (Chapter 2).

An expansion similar to (6) (without estimating the rate of mean-square convergence) was obtained in [4] using a different approach.

The expansion (6) for the narrow particular case  $i_1 = \dots = i_k \neq 0$  can be obtained under the condition of convergence of limiting traces [5] (Theorem 5.1), [6] (Theorem 4.1), [1] (Remark 1.5.7, Proposition 4.1.2) (the definition of limiting traces can be found in [6]).

## REFERENCES

- [1] Budhiraja, A. Multiple stochastic integrals and Hilbert space valued traces with applications to asymptotic statistics and non-linear filtering. Ph. D. Thesis, The University of North Carolina at Chapel Hill, 1994, VII+132 pp.
- [2] Itô, K. Multiple Wiener integral. *J. Math. Soc. Japan*, 3, 1 (1951), 157-169.
- [3] Kuznetsov D.F. Strong Approximation of Iterated Itô and Stratonovich Stochastic Integrals Based on Generalized Multiple Fourier Series. Application to Numerical Solution of Itô SDEs and Semilinear SPDEs. arXiv:2003.14184v28 [math.PR], 2022, 869 pp. DOI: <http://doi.org/10.48550/arXiv.2003.14184>
- [4] Rybakov K.A. Orthogonal expansion of multiple Stratonovich stochastic integrals. *Differencialnie Uravnenia i Protsey Upravlenia*, 4 (2021), 81–115. Available at: <http://diffjournal.spbu.ru/EN/numbers/2021.4/article.1.5.html>
- [5] Johnson, G.W., Kallianpur, G. Homogeneous chaos,  $p$ -forms, scaling and the Feynman integral. *Trans. Amer. Math. Soc.*, 340 (1993), 503-548.
- [6] Budhiraja, A., Kallianpur, G. Two results on multiple Stratonovich integrals. *Statistica Sinica*, 7 (1997), 907-922.