

Shishkova A. A. (Tomsk, Russia) — Hedging problem for the asian option.

Consider a standart Black and Scholes model with several risky assets. We assume that the time horozont is $T = 1$. The riskless asset is a constant over time $B = 1$ and risky assets with price processes $(S_i(t))_{1 \leq i \leq d}$ are driven by the following system of SDEs

$$dS_i(t) = \sigma_i S_i(t) dW_i(t), \quad 0 \leq t \leq 1, \quad i = 1 \dots d.$$

Asian option payoff function is given by $f_1 = (\frac{1}{d} \int_0^1 \sum_{i=1}^d S_i(t) dt - K)_+$, where K – strike price. The main result of the work is the obtained formulas for calculating the hedging strategy $\gamma_i(t) = G'_{y_i}(t, \xi(t), S(t))$, $0 \leq t \leq 1$, $i = 1 \dots d$, where

$$G(t, x, y) = \mathbf{E} \left(\frac{\sum_{i=1}^d x_i + \sum_{i=1}^d y_i \tilde{\eta}_i(v)}{d} - K \right)_+$$

$\xi_i(t) = \int_0^t S_i(v) dv$ and $\tilde{\eta}_i(v) = \int_0^v \exp \{ \sigma_i W_i(u) - \sigma_i^2 u / 2 \}$, $v = 1 - t$. Using the Brownian bridge, we found the densities of random variables $\tilde{\eta}_i(v)$ and studied the analytic properties (differentiability) of the obtained densities. Based on the results presented in [2] we solved the task above. Proved that function $G(t, x, y)$ has continuous derivatives and can be represented by the Ito formula.

REFERENCES

1. *Liptser R.S. and Shiryaev A.N.* Statistics of random processes. 2nd rev. and exp. ed. Springer – Verlag Berlin, 2001, 425 p.
2. Shishkova A.A. Calculation of Asian options for the Black Scholes model. // Bulletin of Tomsk State University. Mathematics and mechanics. 2018. No.51, pp.48–63. DOI: 10.17223/19988621/51/5

This work was supported by the Ministry of Education and Science of the Russian Federation, state assignment No. 2.3208.2017 / 4.6.